

GEOMETRICAL ASPECTS OF EXPANSIONS IN COMPLEX BASES

ANNA CHIARA LAI

ABSTRACT. We study the set of the representable numbers in base $q = pe^{i\frac{2\pi}{n}}$ with $\rho > 1$ and $n \in \mathbb{N}$ and with digits in a arbitrary finite real alphabet A . We give a geometrical description of the convex hull of the representable numbers in base q and alphabet A and an explicit characterization of its extremal points. A characterizing condition for the convexity of the set of representable numbers is also shown.

1. INTRODUCTION

In this paper we deal with expansions with digits in arbitrary alphabets and bases of the form $pe^{i\frac{2\pi}{n}}$ with $p > 1$ and $n \in \mathbb{N}$, namely we are interested in developments in power series of the form

$$(1) \quad \sum_{j=1}^{\infty} \frac{x_j}{q^j}$$

with the coefficients x_j belonging to a finite set of positive real values A named *alphabet* and the ratio $q = pe^{i\frac{2\pi}{n}}$, named *base*. The assumption $p > 1$ ensures the convergence of (1), being the base q greater than 1 in modulus. When a number x satisfies

$$x = \sum_{j=1}^{\infty} \frac{x_j}{q^j},$$

for a sequence $(x_j)_{j \geq 1}$ with digits in the alphabet A , we say that x is *representable* in base q and alphabet A and we call $(x_j)_{j \geq 1}$ a *representation* or *expansion* of x .

The first number systems in complex base seem to be those in base $2i$ with alphabet $\{0, 1, 2, 3\}$ and the one in base $-1 + i$ and alphabet $\{0, 1\}$, respectively introduced by Knuth in [11] and by Penney in [14]. After that many papers were devoted to representability with bases belonging to larger and larger classes of complex numbers, e.g. see [10] for the Gaussian integers in the form $-n \pm i$ with $n \in \mathbb{N}$, [9] for the quadratic fields and [2] for the general case. Loreti and Kormornik pursued the work in [2] by introducing a greedy algorithm for the expansions in complex base with non rational argument [13]. In the eighties a parallel line of research was developed by Gilbert. In [5] he described the fractal nature of the set of the representable numbers, e.g. the set of representable in base $-1 + i$ with digits $\{0, 1\}$ coincides with the fascinating space-filling twin dragon curves [12]. Hausdorff dimension of some set of representable numbers was calculated in [6] and a weaker notion of self-similarity was introduced for the study of the

2000 *Mathematics Subject Classification.* 11A63.

Key words and phrases. Expansions in complex bases, positional numeration systems in complex bases, iterated function systems.

boundary of the representable sets [7]. Complex base numeration systems and in particular the geometry of the set of representable numbers have been widely studied by the point of view of their relations with iterated function systems and tilings of the complex plane, too. For a survey on the topology of the tiles associated to bases belonging to quadratic fields we refer to [1].

We study the convex hull of the set of representable numbers by giving first a geometrical description then an explicit characterization of its extremal points. We also show a characterizing condition for the convexity of the set of representable numbers.

Expansions in complex base have several applications. For example, in the context of computer arithmetics, the interesting property of these numerations systems is that they allow multiplication and division of complex base in a unified manner, without treating real and imaginary part separately — see [12], [6] and [4]. Representation in complex base have been also used in cryptography with the purpose of speeding up onerous computations such as modular exponentiations [3] and multiplications over elliptic curves [17]. Finally we refer to [15] for a dissertation on the applications of the numerations in complex base to the compression of images on fractal tilings.

Organization of the paper. Most of the arguments of this paper laying on geometrical properties, in Section 2 we show some results on complex plane geometry. In Section 3 we characterize the shape and the extremal points of the convex hull of the set of representable numbers. In Section 4 we give a necessary and sufficient condition to have a *convex* set of representable numbers, this property being sufficient for a full representability of complex numbers.

2. GEOMETRICAL BACKGROUND

By using the isometry between $\mathbb{C}$ and $\mathbb{R}^2$ we extend to $\mathbb{C}$ some definitions which are proper of the plane geometry. Elements of $\mathbb{C}$ are considered vectors (or sometimes points) and we endow $\mathbb{C}$ with the scalar product

$$\mathbf{u} \cdot \mathbf{v} := |\mathbf{u}| |\mathbf{v}| \cos(\arg \mathbf{u} - \arg \mathbf{v}).$$

Remark 2.1.

$$\mathbf{u} \cdot \mathbf{v} \leq 0$$

if and only if

$$\cos(\arg \mathbf{u} - \arg \mathbf{v}) \leq 0.$$

A polygon in the complex plane is the bounded region of $\mathbb{C}$ contained in a closed chain of segments, the *edges*, whose endpoints are the *vertices*. If two adjacent edges belong to the same line, namely if they are adjacent and parallel, then they are called *consecutive* and their common endpoint is a *degenerate vertex*. If a vertex is not degenerate, it is called *extremal point*. The index operations on the vertices $\mathbf{v}_1, \dots, \mathbf{v}_H \in \mathbb{C}$ of a polygon are always considered modulus their number, namely

$$\mathbf{v}_{h+s} := \mathbf{v}_{h+s \bmod H}$$

so that for instance $\mathbf{v}_0 = \mathbf{v}_H$ and $\mathbf{v}_{H+1} = \mathbf{v}_1$.

Consider an edge whose endpoints are two vertices $\mathbf{v}_{h-1}$ and $\mathbf{v}_h$ and its *normal vector*

$$(2) \quad \mathbf{n}_h = (\mathbf{v}_h - \mathbf{v}_{h-1})^\perp := -i(\mathbf{v}_h - \mathbf{v}_{h-1})$$

A set of vertices $\{\mathbf{v}_1, \dots, \mathbf{v}_H\}$ is *counter-clockwise ordered* if there exists $s \in \{0, \dots, H-1\}$ such that for every $h = 1, \dots, H-1$

$$(3) \quad \arg \mathbf{n}_{h+s} \leq \arg \mathbf{n}_{h+s+1}.$$

A set of vertices $\{\mathbf{v}_1, \dots, \mathbf{v}_H\}$ is *clockwise ordered* if there exists $s \in \{0, \dots, H-1\}$ such that for every $i = 1, \dots, H-1$

$$(4) \quad \arg \mathbf{n}_{h+s} \geq \arg \mathbf{n}_{h+s+1}.$$

Remark 2.2. A polygon is convex if and only if its vertices are clockwise or counter-clockwise ordered (see [8] for the $\mathbb{R}^2$ case, the complex case readily follows by employing the isometry $(x, y) \mapsto x + iy$).

We are now interested in establishing condition on a point $x \in \mathbb{C}$ to belong to a convex polygon. The following result is an adapted version of the exterior criterion for the Point-In-Polygon problem (see for instance [16]).

Proposition 2.3. Let P be a convex polygon whose vertices are $\mathbf{v}_1, \dots, \mathbf{v}_H$. Then a complex value x belongs to P if and only if for every $h = 1, \dots, H$

$$(x - \mathbf{v}_h) \cdot \mathbf{n}_h \leq 0.$$

The *convex hull* of a set $X \subset \mathbb{C}$ is the smallest convex set containing X and it is denoted by using the symbol $\text{conv}(X)$. When X is finite, its convex hull is a convex polygon whose vertices are in X .

We now study the convex hull of the set $P \cup (P + \mathbf{t})$, being P a convex polygon, $\mathbf{t} \in \mathbb{C}$ and $P + \mathbf{t} := \{x + \mathbf{t} \mid x \in P\}$.

Remark 2.4. If $\mathbf{t} \in \mathbb{C}$ then

$$\arg \mathbf{t}^\perp = \left(\arg \mathbf{t} + \frac{3\pi}{2} \right) \bmod 2\pi$$

and

$$\arg(-\mathbf{t})^\perp = \left(\arg \mathbf{t} + \frac{\pi}{2} \right) \bmod 2\pi$$

Theorem 2.5. Let P be a convex polygon with counter-clockwise ordered vertices in $\{\mathbf{v}_1, \dots, \mathbf{v}_H\}$ and let $\mathbf{t} \in \mathbb{C}$. Denote h_1 and h_2 the indices respectively satisfying

$$(5) \quad \arg \mathbf{n}_{h_1} \leq \arg -\mathbf{t}^\perp < \arg \mathbf{n}_{h_1+1}$$

$$(6) \quad \arg \mathbf{n}_{h_2} \leq \arg \mathbf{t}^\perp < \arg \mathbf{n}_{h_2+1}$$

Then the convex hull of $P \cup (P + \mathbf{t})$ is a polygon whose vertices are:

$$(7) \quad \mathbf{v}_{h_1}, \dots, \mathbf{v}_{h_2}, \mathbf{v}_{h_2} + \mathbf{t}, \dots, \mathbf{v}_{h_1-1} + \mathbf{t}, \mathbf{v}_{h_1} + \mathbf{t}.$$

Remark 2.6. As the index operations on the vertices are considered modulus H , if $h_1 > h_2$ the expression in (7) means

$$(8) \quad \mathbf{v}_{h_1}, \dots, \mathbf{v}_H, \mathbf{v}_1, \dots, \mathbf{v}_{h_2}, \mathbf{v}_{h_2} + \mathbf{t}, \dots, \mathbf{v}_{h_1-1} + \mathbf{t}, \mathbf{v}_{h_1} + \mathbf{t}.$$

conversely if $h_1 < h_2$ the extended version of (7) is

$$(9) \quad \mathbf{v}_{h_1}, \dots, \mathbf{v}_{h_2}, \mathbf{v}_{h_2} + \mathbf{t}, \dots, \mathbf{v}_H + \mathbf{t}, \mathbf{v}_1 + \mathbf{t}, \dots, \mathbf{v}_{h_1-1} + \mathbf{t}, \mathbf{v}_{h_1} + \mathbf{t}.$$

Proof. As P is a convex polygon, its vertices are either clockwise or counter-clockwise ordered. We may assume without loss of generality the latter case and, in particular, that

$$(10) \quad \arg(\mathbf{n}_h) \leq \arg(\mathbf{n}_{h+1})$$

for every $h = 1, \dots, H-1$. Therefore h_1 and h_2 are well defined. By definition $h_1 \neq h_2$, hence we may distinguish the cases $h_1 > h_2$ and $h_1 < h_2$. We discuss only the latter case, because the proves are similar. We use the symbol $\bar{P}$ to denote the polygon whose vertices are listed in (7), in particular we set

$$\bar{\mathbf{v}}_h = \begin{cases} \mathbf{v}_{h+h_1-1} & \text{if } h = 1, \dots, h_2 - h_1 + 1 \\ \mathbf{v}_{h+h_1-2} + \mathbf{t} & \text{if } h = h_2 - h_1 + 2, \dots, H + 2 \end{cases}$$

Remark that by shifting the vertices of $\bar{P}$ of $s := H - h_1 + 2$ positions, we obtain the ordered list

$$\mathbf{v}_1 + \mathbf{t}, \dots, \mathbf{v}_{h_1} + \mathbf{t}, \mathbf{v}_{h_1}, \dots, \mathbf{v}_{h_2}, \mathbf{v}_{h_2} + \mathbf{t}, \dots, \mathbf{v}_H + \mathbf{t}$$

and, in particular,

$$\bar{\mathbf{v}}_{h+s} = \begin{cases} \mathbf{v}_h + \mathbf{t} & \text{if } h = 1, \dots, h_1 \\ \mathbf{v}_{h-1} & \text{if } h = h_1 + 1, \dots, h_2 + 1 \\ \mathbf{v}_{h-2} + \mathbf{t} & \text{if } h = h_2 + 2, \dots, H + 2. \end{cases}$$

Hence

$$\bar{\mathbf{n}}_{h+s} = \begin{cases} \mathbf{n}_h & \text{if } h = 1, \dots, h_1 \\ (-\mathbf{t})^\perp & \text{if } h = h_1 + 1 \\ \mathbf{n}_{h-1} & \text{if } h = h_1 + 2, \dots, h_2 + 1 \\ (\mathbf{t})^\perp & \text{if } h = h_2 + 2 \\ \mathbf{n}_{h-2} & \text{if } h = h_2 + 3, \dots, H + 2. \end{cases}$$

Thus the definition of h_1 and of h_2 , together with (10), implies that for every $h = 1, \dots, H+1$

$$(11) \quad \arg \bar{\mathbf{n}}_{h+s} \leq \arg \bar{\mathbf{n}}_{h+s+1}$$

namely the vertices of $\bar{P}$ are counter-clockwise ordered. In view of Remark 2.2 we may deduce that $\bar{P}$ is convex and, in particular,

$$\bar{P} = \text{conv}\{\bar{\mathbf{v}}_h \mid h = 1, \dots, H + 2\}.$$

Now we want to show

$$\bar{P} = \text{conv}(P \cup (P + \mathbf{t}))$$

by double inclusion. Since P is convex, we have

$$P = \text{conv}\{\mathbf{v}_h \mid h = 1, \dots, H\},$$

$$P + \mathbf{t} = \text{conv}\{\mathbf{v}_h + \mathbf{t} \mid h = 1, \dots, H\}$$

and

$$\text{conv}(P \cup (P + \mathbf{t})) = \text{conv}\{\mathbf{v}_h, \mathbf{v}_h + \mathbf{t} \mid h = 1, \dots, H\},$$

therefore

$$\bar{P} = \text{conv}\{\bar{\mathbf{v}}_h \mid h = 1, \dots, H\} \subseteq \text{conv}(P \cup (P + \mathbf{t})).$$

To prove the other inclusion, it suffices to show that $P \cup (P + \mathbf{t}) \subseteq \bar{P}$. In view of Proposition 2.3, this is equivalent to prove that for every $y \in P \cup (P + \mathbf{t})$ and for every $h = 1, \dots, 2H$

$$(12) \quad (y - \bar{\mathbf{v}}_h) \cdot \bar{\mathbf{n}}_h \leq 0.$$

Now consider $y \in P \cup (P + \mathbf{t})$ and remark that

$$y = x + \alpha \mathbf{t}$$

for some $x \in P$ and $\alpha \in \{0, 1\}$. Therefore we may rewrite (12) as follows

$$(13) \quad (x + \alpha \mathbf{t} - \bar{\mathbf{v}}_h) \cdot \bar{\mathbf{n}}_h \leq 0$$

for every $h = 1, \dots, H + 2$. First of all remark that, by Proposition 2.3, for every $h = 1, \dots, H$

$$(14) \quad (x - \mathbf{v}_h) \cdot \mathbf{n}_h \leq 0$$

and, consequently,

$$(15) \quad \begin{aligned} (x - \mathbf{v}_h) \cdot \mathbf{n}_{h+1} &= (x - \mathbf{v}_{h+1}) \cdot \mathbf{n}_{h+1} + (\mathbf{v}_{h+1} - \mathbf{v}_h) \cdot \mathbf{n}_{h+1} \\ &= (x - \mathbf{v}_{h+1}) \cdot \mathbf{n}_{h+1} + (\mathbf{v}_{h+1} - \mathbf{v}_h) \cdot (\mathbf{v}_{h+1} - \mathbf{v}_h)^\perp \\ &= (x - \mathbf{v}_{h+1}) \cdot \mathbf{n}_{h+1} \\ &\leq 0; \end{aligned}$$

Therefore,

$$(16) \quad \max\{\cos(\arg(x - \mathbf{v}_h) - \arg \mathbf{n}_h), \cos(\arg(x - \mathbf{v}_h) - \arg \mathbf{n}_{h+1})\} \leq 0.$$

Now,

- if $h = 1$ then the definition of h_1 , namely

$$\arg \mathbf{n}_{h_1} \leq \arg(-\mathbf{t})^\perp < \arg \mathbf{n}_{h_1+1},$$

and (16) imply

$$\begin{aligned} \cos(\arg(x - \mathbf{v}_h) - \arg(-\mathbf{t})^\perp) \\ \leq \max\{\cos(\arg(x - \mathbf{v}_{h_1}) - \arg \mathbf{n}_{h_1}), \cos(\arg(x - \mathbf{v}_{h_1}) - \arg \mathbf{n}_{h_1+1})\} \\ \leq 0. \end{aligned}$$

Then

$$(x + \alpha \mathbf{t} - \bar{\mathbf{v}}_1) \cdot \bar{\mathbf{n}}_1 = (x - \mathbf{v}_{h_1}) \cdot (-\mathbf{t})^\perp + \alpha \mathbf{t} \cdot (-\mathbf{t})^\perp \leq 0.$$

- If $h = 2, \dots, h_2 - h_1$ then

$$(x + \alpha \mathbf{t} - \bar{\mathbf{v}}_h) \cdot \bar{\mathbf{n}}_h = (x - \mathbf{v}_{h+h_1-1}) \cdot \mathbf{n}_{h+h_1-1} + \alpha \mathbf{t} \cdot \mathbf{n}_{h+h_1-1} \leq 0,$$

because $\mathbf{t} \cdot \mathbf{n}_h \leq 0$ whenever $h_1 \leq h \leq h_2$. Indeed

$$\left(\arg \mathbf{t} + \frac{\pi}{2}\right) \bmod 2\pi = \arg(-\mathbf{t}^\perp) \leq \arg \mathbf{n}_h \leq \arg \mathbf{t}^\perp = \left(\arg \mathbf{t} + \frac{3\pi}{2}\right) \bmod 2\pi$$

hence

$$\cos(\arg \mathbf{t} - \arg \mathbf{n}_h) \leq \max\left\{\cos\left(\frac{\pi}{2}\right), \cos\left(\frac{3\pi}{2}\right)\right\} = 0.$$

- If $h = h_2 - h_1 + 1$ then

$$(x + \alpha \mathbf{t} - \bar{\mathbf{v}}_h) \cdot \bar{\mathbf{n}}_h = (x + \alpha \mathbf{t} - \mathbf{v}_{h_2} - \mathbf{t}) \cdot \mathbf{t}^\perp = (x - \mathbf{v}_{h_2}) \cdot \mathbf{t}^\perp \leq 0$$

Indeed

$$\arg \mathbf{n}_{h_2} \leq \arg \mathbf{t}^\perp < \arg \mathbf{n}_{h_2+1},$$

and (16) imply

$$\begin{aligned} \cos(\arg(x - \mathbf{v}_h) - \arg(\mathbf{t})^\perp) &\leq \\ &\leq \max\{\cos(\arg(x - \mathbf{v}_{h_2}) - \arg \mathbf{n}_{h_2}), \cos(\arg(x - \mathbf{v}_{h_2}) - \arg \mathbf{n}_{h_2+1})\} \\ &\leq 0. \end{aligned}$$

- Finally if $h = h_2 - h_1 + 2, \dots, H + 2$ then

$$\begin{aligned} (x + \alpha \mathbf{t} - \bar{\mathbf{v}}_h) \cdot \bar{\mathbf{n}}_h &= (x + \alpha \mathbf{t} - \mathbf{v}_{h+h_1-2} - \mathbf{t}) \cdot \mathbf{n}_{h_1-2+h} \\ &\leq (x - \mathbf{v}_{h+h_1-2}) \cdot \mathbf{n}_{h_1-2+h} - (1 - \alpha) \mathbf{t} \cdot \mathbf{n}_{h_1-2+h} \\ &\leq 0 \end{aligned}$$

because $1 - \alpha \geq 0$ and $\mathbf{t} \cdot \mathbf{n}_{h_1-2+h} \geq 0$ for every $h = h_2 - h_1 + 2, \dots, H + 2$.

Hence (13) holds for every $x \in P$ and $\alpha \in \{0, 1\}$ and the proof is complete. $\square$

Corollary 2.7. *Let P be a convex polygon with l edges and let $\mathbf{t} \in \mathbb{C}$. Then $\text{conv}(P \cup (P + \mathbf{t}))$ has $H + 2$ (possibly consecutive) edges, in particular l edges of $\text{conv}(P \cup (P + \mathbf{t}))$ are parallel to the edges of P and 2 edges are parallel to $\mathbf{t}$.*

Proof. It immediately follows by the list of vertices given in (7). $\square$

Corollary 2.8. *Let P be a convex polygon with e extremal points and let $\mathbf{t} \in \mathbb{C}$. Then:*

- (a) *if $\mathbf{t}$ is not parallel to any edge of P then $\text{conv}(P \cup (P + \mathbf{t}))$ has $e + 2$ extremal points;*
- (b) *if $\mathbf{t}$ is parallel to 1 edge of P then $\text{conv}(P \cup (P + \mathbf{t}))$ has $e + 1$ extremal points;*
- (c) *if $\mathbf{t}$ is parallel to 2 edges of P then $\text{conv}(P \cup (P + \mathbf{t}))$ has e extremal points.*

Proof. Set l the number of edges of P and $d := l - e$ the number of degenerate vertices. By definition, a vertex $\mathbf{v}$ is degenerate if the normal vectors of its adjacent edges have the same argument. It follows by (7) that the normal vectors of $\text{conv}(P \cup (P + \mathbf{t}))$ are the following:

$$\mathbf{n}_{h_1}, \dots, \mathbf{n}_{h_2-1}, \mathbf{t}^\perp, \mathbf{n}_{h_2}, \dots, \mathbf{n}_{h_1-1}, -\mathbf{t}^\perp$$

with h_1 and h_2 satisfying:

$$\begin{aligned} \arg \mathbf{n}_{h_1} &\leq \arg -\mathbf{t}^\perp < \arg \mathbf{n}_{h_1+1} \\ \arg \mathbf{n}_{h_2} &\leq \arg \mathbf{t}^\perp < \arg \mathbf{n}_{h_2+1} \end{aligned}$$

Thus by denoting l_t the number of vertices of P_t , by d_t the number of the degenerate vertices and by e_t the number of extremal points, by Corollary 2.7 we have that $l_t = l + 2$ and :

$$d_t = \begin{cases} d & \text{if } \mathbf{t}^\perp \text{ is neither parallel to } \mathbf{n}_{h_1} \text{ nor to } \mathbf{n}_{h_2-1}; \\ d + 1 & \text{if } \mathbf{t}^\perp \text{ is parallel to } \mathbf{n}_{h_1} \text{ or to } \mathbf{n}_{h_2-1}; \\ d + 2 & \text{if } \mathbf{t}^\perp \text{ is parallel to } \mathbf{n}_{h_1} \text{ and to } \mathbf{n}_{h_2-1}. \end{cases}$$

Hence thesis follows by the relation $e_t = l_t - d_t$. $\square$

We conclude this section with the following result on the convexity of $P \cup P + \mathbf{t}$.

Proposition 2.9. *Let P be a convex polygon and assume that 2 edges of P are parallel to the real axis on the complex plane and that their length is equal to 1. Consider $t_1, \dots, t_m \in \mathbb{R}$ such that*

$$(17) \quad t_1 < \dots < t_m.$$

Then

$$\max_{i=1, \dots, m-1} t_{i+1} - t_i \leq 1$$

if and only if

$$\bigcup_{i=1}^m (P + t_i)$$

is a convex set.

Proof. Only if part.

Suppose $t_{i_1+1} - t_{i_1} > 1$ for some $i_1 \in \{1, \dots, m-1\}$. Let $\mathbf{v}_1$ and $\mathbf{v}_2$ two extremal points of P such that

$$\operatorname{re}(\mathbf{v}_1) \leq \operatorname{re}(\mathbf{v}_2)$$

and

$$\operatorname{im}(\mathbf{v}_1) = \operatorname{im}(\mathbf{v}_2)$$

By the inequality above, the edge with endpoints $\mathbf{v}_1$ and $\mathbf{v}_2$ is parallel to the real axis and, consequently,

$$(18) \quad \operatorname{re}(\mathbf{v}_2) - \operatorname{re}(\mathbf{v}_1) \leq 1 < t_{i_1+1} - t_{i_1}.$$

Now define for every $i = 1, \dots, m$

$$\mathbf{e}_i := \{\mathbf{v}_1 + x \mid x \in [t_i, t_i + \operatorname{re} \mathbf{v}_2 - \operatorname{re} \mathbf{v}_1]\}$$

and remark that the convexity of P implies

$$\bigcup_{i=1}^m P + t_i \cap \{\mathbf{v}_1 + t \mid t \in \mathbb{R}\} = \bigcup_{i=1}^m \mathbf{e}_i$$

By (17), for every $\alpha \in (0, 1)$ and every $i = 1, \dots, m-1$

$$\alpha(\mathbf{v}_2 + t_i) + (1 - \alpha)(\mathbf{v}_1 + t_{i+1}) \in \bigcup_{i=1}^m \mathbf{e}_i$$

if and only if

$$\alpha(\mathbf{v}_2 + t_i) + (1 - \alpha)(\mathbf{v}_1 + t_{i+1}) \in \mathbf{e}_i \cap \mathbf{e}_{i+1}.$$

Now, setting $\mathbf{x}_1 := \mathbf{v}_2 + t_{i_1}$ and $\mathbf{x}_2 := \mathbf{v}_1 + t_{i_1+1}$, we have that for every $\alpha \in (0, 1)$

$$\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2 \in \operatorname{conv} \left(\bigcup_{i=1}^m (P + t_i) \right)$$

and

$$\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2 \in \{\mathbf{v}_1 + t \mid t \in \mathbb{R}\}$$

but, in view of (18),

$$\mathbf{e}_{i_1} \cap \mathbf{e}_{i_1+1} = \emptyset$$

hence

$$\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2 \notin \bigcup_{i=1}^m (P + t_i).$$

Therefore $\bigcup_{i=1}^m (P + t_i)$ is not a convex set.

If part

Let m_P and M_P be such that P is a subset of $\{x \in \mathbb{C} \mid m_P \leq \text{im}(x) \leq M_P\}$. For every $m_P \leq y \leq M_P$ we consider the set

$$I_y := \{x \in \mathbb{R} \mid x + iy \in P\}$$

As P is convex, I_y is an interval, whose endpoints are denoted by a_y and b_y . The convexity of P also implies $|I_y| \geq 1$; therefore $t_{i+1} - t_i \leq 1$ implies

$$[a_y + t_1, b_y + t_m] = \bigcup_{i=1}^m [a_y + t_i, b_y + t_i].$$

and

$$(19) \quad x + iy \in \bigcup_{i=1}^m (P + t_i) \quad \text{if and only if} \quad x \in [a_y + t_1, b_y + t_m].$$

We want to prove that $\bigcup_{i=1}^m (P + t_i)$ is a convex set by showing that it contains any convex combination of its points. So fix

$$\mathbf{x}_1, \mathbf{x}_2 \in \bigcup_{i=1}^m (P + t_i).$$

If $\mathbf{x}_1$ and $\mathbf{x}_2$ are both in $P + t_i$ for some $i = 1, \dots, m$ the convexity of P implies the thesis. Otherwise suppose $\mathbf{x}_1 \in P + t_{i_1}$ and $\mathbf{x}_2 \in P + t_{i_2}$ with $t_{i_1} < t_{i_2}$ and consider a convex combination $\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2$, with $\alpha \in [0, 1]$. Remark that $\mathbf{x}_2 \in P + t_{i_2}$ implies $\mathbf{x}_2 - (t_{i_2} - t_{i_1}) \in P + t_{i_1}$ and, consequently,

$$\mathbf{x} := \alpha \mathbf{x}_1 + (1 - \alpha)(\alpha \mathbf{x}_2 - \mathbf{t}) \in P + t_{i_1}.$$

Therefore

$$\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2 = \mathbf{x} + (1 - \alpha) \mathbf{t} = x + iy + (1 - \alpha)(t_{i_2} - t_{i_1}).$$

for some $x \in I_y + t_{i_1}$ and $y \in [m_P, M_P]$. Hence

$$\begin{aligned} a_y + t_1 &\leq a_y + t_{i_1} \leq x + (1 - \alpha)(t_{i_2} - t_{i_1}) \leq b_y + t_{i_1} + (1 - \alpha)(t_{i_2} - t_{i_1}) \\ &\leq b_y + t_m. \end{aligned}$$

In view of (19) we finally get $\alpha \mathbf{x}_1 + (1 - \alpha) \mathbf{x}_2 \in \bigcup_{i=1}^m (P + t_i)$ and hence the thesis. $\square$

3. CHARACTERIZATION OF THE CONVEX HULL OF REPRESENTABLE NUMBERS

In this section we investigate the shape of the convex hull of the set of representable numbers in base $pe^{\frac{2\pi}{n}i}$ and with alphabet A . We adopt the following notations.

Notation 3.1. We use the symbol $\Lambda_{n,p,A}$ to denote the set of representable numbers in base $q_{n,p} := pe^{\frac{2\pi}{n}i}$ and with alphabet A , namely

$$\Lambda_{n,p,A} = \left\{ \sum_{j=1}^{\infty} \frac{x_j}{q_{n,p}^j} \mid x_j \in A \right\}.$$

We set

$$X_{n,p} := \left\{ \sum_{k=0}^{n-1} x_k q_{n,p}^k \mid x_k \in \{0, 1\} \right\}$$

and

$$P_{n,p} := \text{conv}(X_{n,p}).$$

Remark 3.2. As $X_{n,p}$ is finite, $P_{n,p}$ is a polygon.

The following result represents a first simplification of our problem: in fact the characterization of the convex hull of the infinite set $\Lambda_{n,p,A}$ is showed to be equivalent to the study of $P_{n,p}$.

Lemma 3.3 (Farkas' Lemma). *Let $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^n$. The system $A\mu = b$ admits a non-negative solution if and only if for every $u \in \mathbb{R}^m$ the inequality $A^\top \cdot u \geq 0$ implies $b^\top \cdot u \geq 0$.*

Lemma 3.4. *Let $q \in \mathbb{C}$, $n \in \mathbb{N}$ and $\lambda_k \in [\lambda_{\min}, \lambda_{\max}]$ for every $k = 0, \dots, n-1$, $\lambda_{\min}, \lambda_{\max} \in \mathbb{R}$ and*

$$S := \{(x_0^{(h)}, \dots, x_{n-1}^{(h)}) \mid x_k^{(h)} \in \{\lambda_{\min}, \lambda_{\max}\}; k = 1, \dots, n; h = 1, \dots, 2^n\}$$

be the set of sequences of length k and with digits in $\{\lambda_{\min}, \lambda_{\max}\}$.

Then there exist $\mu_1, \dots, \mu_{2^n} \geq 0$ such that

$$\sum_{h=1}^{2^n} \mu_h = 1$$

and

$$(20) \quad \sum_{k=0}^{n-1} q^k \lambda_k = \sum_{h=1}^{2^n} \mu_h \sum_{k=0}^{n-1} q^k x_k^{(h)}$$

Proof. Consider the linear system with 2^n indeterminates $\mu_1, \dots, \mu_{2^n}$ and with $n+1$ equations

$$\begin{cases} \sum_{h=1}^{2^n} \mu_h x_k^{(h)} = \lambda_k & \text{for } k = 0, \dots, n-1; \\ \sum_{h=1}^{2^n} \mu_h = 1. \end{cases}$$

We may rewrite the above system in the form

$$A\mu = b$$

where the h -th column of A satisfies $A_h^\top = (x_1^{(h)}, \dots, x_k^{(h)}, 1)$ and $b^\top = (\lambda_1, \dots, \lambda_k, 1)$. In order to apply Farkas' Lemma consider $u \in \mathbb{R}^{n+1}$ such that

$$A_h^\top \cdot u = \sum_{k=1}^n x_k^{(h)} u_k + u_{n+1} \geq 0$$

for every $h = 1, \dots, 2^n$. We recall that $\lambda_k \in [\lambda_{\min}, \lambda_{\max}]$ for every k , then we may consider the sequence $(x_1^{(h)}, \dots, x_n^{(h)}) \in S$ such that for every $k = 1, \dots, n$

$$x_k^{(h)} \geq \lambda_k \quad \text{if and only if} \quad u_k \geq 0.$$

Consequently

$$\sum_{k=1}^n (\lambda_k - x_k^{(h)}) u_k \geq 0$$

and

$$b^\top \cdot u = \sum_{k=1}^n \lambda_k u_k + u_{n+1} \geq \sum_{k=1}^n x_k^{(h)} u_k + u_{n+1} \geq 0.$$

By Farkas' Lemma, there exist $\mu_1, \dots, \mu_{2^n} \geq 0$ with $\sum_{h=1}^{2^n} \mu_h = 1$ such that

$$\sum_{h=1}^{2^n} \mu_h x_k^{(h)} = \lambda_k;$$

for every $k = 1, \dots, n$ and, consequently, $\mu_1, \dots, \mu_{2^n}$ also satisfy

$$\sum_{k=0}^{n-1} q^k \lambda_k = \sum_{k=0}^{n-1} q^k \sum_{h=1}^{2^n} \mu_h x_k^{(h)} = \sum_{h=1}^{2^n} \mu_h \sum_{k=0}^{n-1} q^k x_k^{(h)}.$$

□

Proposition 3.5. For every $n \geq 1$, $p > 1$ and $q_{n,p} = pe^{\frac{2\pi}{n}i}$:

$$(21) \quad \text{conv}(\Lambda_{n,p,A}) = \frac{\max A - \min A}{p^n - 1} \cdot P_{n,p} + \frac{1}{p^n - 1} \sum_{k=0}^{n-1} \min A \, q_{n,p}^k.$$

Proof. Fix n and p and, in order to lighten the notations, set $q = q_{n,p}$, $X_n = X_{n,p}$ and $P_n = P_{n,p}$. Consider

$$x = \sum_{j=1}^{\infty} \frac{x_j}{q^j} \in \Lambda_{n,p,A}.$$

As $q^n = p^n$,

$$(22) \quad \begin{aligned} \sum_{j=1}^{\infty} \frac{x_j}{q^j} &= \sum_{k=0}^{n-1} q^k \sum_{j=1}^{\infty} \frac{x_{jn-k}}{p^{jn}} \\ &= \sum_{k=0}^{n-1} q^k \lambda_k \end{aligned}$$

with

$$\lambda_k := \sum_{j=1}^{\infty} \frac{x_{jn-k}}{p^{jn}} \in \left[\frac{\min A}{p^n - 1}, \frac{\max A}{p^n - 1} \right].$$

Therefore by Lemma 3.4 any element of $\Lambda_{n,p,A}$ is a convex combination of complex numbers that can be written in the form

$$\sum_{k=0}^{n-1} q^k \tilde{x}_k$$

with $\tilde{x}_k \in \left\{ \frac{\min A}{p^n - 1}, \frac{\max A}{p^n - 1} \right\}$. Thus

$$\begin{aligned} \text{conv}(\Lambda_{n,p,A}) &= \text{conv} \left(\left\{ \sum_{k=0}^{n-1} q^k \tilde{x}_k \mid \tilde{x}_k \in \left\{ \frac{\min A}{p^n - 1}, \frac{\max A}{p^n - 1} \right\} \right\} \right) \\ &= \frac{\max A - \min A}{p^n - 1} \text{conv}(X_{n,p}) + \frac{1}{p^n - 1} \sum_{k=0}^{n-1} \min A q^k. \end{aligned}$$

□

We now give a geometrical description of $\text{conv}(\Lambda_{n,p,A})$.

Theorem 3.6 (Convex hull of the representable numbers in complex base). *For every $n \geq 1$, $p > 1$ and alphabet A , the set $\text{conv}(\Lambda_{n,p,A})$ is a polygon with the following properties:*

- (a) *the edges are pairwise parallel to $q^0, \dots, q^{n-1}$, where $q = pe^{\frac{2\pi i}{n}}$;*
- (b) *if n is odd then $\text{conv}(\Lambda_{n,p,A})$ has $2n$ extremal points;*
- (c) *if n is even then $\text{conv}(\Lambda_{n,p,A})$ has n extremal points.*

Proof. Fix n and p and, in order to lighten the notations, set $q = q_{n,p}$, $X_n = X_{n,p}$ and $P_n = P_{n,p}$. Our proof is based on showing P_n to have properties (a), (b) and (c); indeed these properties are invariant by rescaling and translation and, by Proposition 3.5, they extend from P_n to $\text{conv}(\Lambda_{n,p,A})$.

We divide the proof in three parts.

Part 1. The edges of P_n are pairwise parallel to $q^0, \dots, q^{n-1}$.

To the end of studying P_n , we consider the sets

$$X_m := \left\{ \sum_{j=0}^{m-1} x_j q^j \mid x_j \in \{0, 1\} \right\};$$

for $m = 1, \dots, n$. We have

$$(23) \quad \begin{cases} X_1 = \{0, 1\}; \\ X_m = X_{m-1} \cup (X_{m-1} + q^{m-1}) \end{cases}$$

and consequently

$$(24) \quad \begin{cases} P_1 = [0, 1]; \\ P_m = \text{conv}(P_{m-1} \cup (P_{m-1} + q^{m-1})). \end{cases}$$

We remark that P_1 can be looked at as a polygon with two vertices and with two overlapped edges that are parallel to q^0 . By iteratively applying Corollary 2.7 we deduce that P_n has pairwise parallel edges and every couple of edges is either parallel to q^0 or to any of the successive translation, i.e. $q^1, \dots, q^{n-1}$.

Part 2. If n is odd then P_n has $2n$ extremal points.

First remark that n odd implies that q^j and q^k are not parallel for every $j \neq k$. We showed above that for every $m = 1, \dots, n$ the edges of the polygon P_{m-1} are parallel to $q^0, \dots, q^{m-2}$ and, consequently, the translation q^{m-1} is not parallel to any edge. Hence, denoting e_m the number of the extremal points of P_m , the first part of Corollary 2.7 implies that e_m is defined by the recursive relation:

$$\begin{cases} e_1 = 2 \\ e_m = e_{m-1} + 2 \end{cases}$$

for every $m = 1, \dots, n$. Hence $e_n = 2n$.

Part 3. If n is even then P_n has n extremal points.

If n is even then $q_n^{m+n/2}$ is parallel to q_n^m for every $m = 1, \dots, n/2$. Since P_m has pairwise parallel edges, we deduce by (a) and (c) in Corollary 2.8 that e_m is defined by the relation:

$$\begin{cases} e_0 = 0; \\ e_m = e_{m-1} + 2 & \text{if } m = 1, \dots, n/2; \\ e_m = e_{m-1} & \text{if } m = n/2 + 1, \dots, n; \end{cases}$$

hence $e_n = n$ and this concludes the proof. $\square$

Example 3.7. *If $n = 3$ and if $p > 1$ then $\text{conv}(\Lambda_{n,p,A})$ is an hexagon.*

If $n = 4$ and if $p > 1$ then $\text{conv}(\Lambda_{n,p,A})$ is a rectangle.




FIGURE 1. Convex hulls of $X_{3,2^{1/3}}$ and of $X_{4,2^{1/4}}$. Remark that when $A = \{0, 1\}$ then $P_{n,p} = \text{conv}(X_{n,p})$ coincides with $\frac{1}{p^n - 1} \text{conv}(\Lambda_{n,p,A})$.

After establishing the shape of $\text{conv}(\Lambda_{n,p,A})$, we are now interested on the explicit characterization of its extremal points. By Proposition 3.5, this is equivalent to characterize the extremal points of $P_{n,p}$ and we shall focus on this problem. Let us see some examples.




FIGURE 2. The sets $\Lambda_{n,2^{1/2},\{0,1\}}$ with $n = 3, \dots, 8$ and their convex hulls. $\Lambda_{n,2^{1/2},\{0,1\}}$ is approximated with the set of expansions with length 14. Remark that $q_{8,2^{1/2}} = 1 + i$ is a Gaussian integer that has been studied, for instance, in [5].

Example 3.8. Set $p > 1$ and $q := q_{3,p} = pe^{\frac{2\pi}{3}i}$. By a direct computation, the set of extremal points of $P_{3,p}$, say $\mathcal{E}(P_{3,p})$, is

$$\mathcal{E}(P_{3,p}) = \{1, 1 + q, q, q + q^2, q^2, q^2 + 1\}.$$

Example 3.9. Set $p > 1$ and $q := q_{4,p} = pe^{\frac{2\pi}{4}i}$. By a direct computation, the set of extremal points of $P_{4,p}$, say $\mathcal{E}(P_{4,p})$, is

$$\mathcal{E}(P_{4,p}) = \{1 + q, q + q^2, q^2 + q^3, q^3 + 1\}.$$

Example 3.8 and Example 3.9 suggest that the set of extremal points of $P_{n,p}$ has an internal structure. In general the vertices of $P_{n,p}$ are element of $X_{n,p}$ and in particular they are (finite) expansions in base $q_{n,p}$. In order to make the structure




FIGURE 3. Extremal points of $P_{3,p}$ with $p = 2^{1/3}$. When the alphabet is $\{0, 1\}$, this coincides with the convex hull of $\Lambda_{A,n,p}$.

of the extremal points more evident, in next examples we focus on the sequences of binary coefficients associated to the extremal points. This point of view requires some notations.

Notation 3.10. Let $(x_0 \cdots x_{n-1})$ be a sequence in $\{0, 1\}^n$ and $q \in \mathbb{C}$. We define

$$(x_0 \cdots x_{n-1})_q := \sum_{k=0}^{n-1} x_k q^k$$

and we introduce σ , the circular shift on finite sequences:

$$\sigma(x_0 x_1 \cdots x_{n-1}) := (x_1 \cdots x_{n-1} x_0).$$

The closure of $(x_0 \cdots x_{n-1})$ with respect to σ is denoted by

$$\text{Orb}(x_0 \cdots x_{n-1}) := \{\sigma^h(x_0 x_1 \cdots x_{n-1}) \mid h = 0, \dots, n-1\}.$$

Finally we define

$$\text{Orb}(x_0 \cdots x_{n-1})_q := \{\sigma^h(x_0 x_1 \cdots x_{n-1})_q \mid h = 0, \dots, n-1\}.$$

Example 3.11. Following relations hold for every $p > 1$ and they are established by means of a symbolic computer program.

If $q = q_{3,p}$ then

$$\begin{aligned} \mathcal{E}(P_{3,p}) &= \{1, 1+q, q, q+q^2, q^2, q^2+1\}. \\ &= \{(100)_q, (110)_q, (010)_q, (011)_q, (001)_q, (101)_q\} \\ &= \text{Orb}(100)_q \cup \text{Orb}(110)_q; \end{aligned}$$

If $q = q_{4,p}$ then

$$\begin{aligned}\mathcal{E}(P_{4,p}) &= \{1 + q, q + q^2, q^2 + q^3, q^3 + 1\}. \\ &= \{(1100)_q, (0110)_q, (0011)_q, (1001)_q\} \\ &= \text{Orb}(1100)_q;\end{aligned}$$

If $q = q_{5,p}$ then

$$\begin{aligned}\mathcal{E}(P_{5,p}) &= \{(11000)_q, (11100)_q, (01100)_q, (01110)_q, (00110)_q, \\ &\quad (00111)_q, (00011)_q, (10011)_q, (10001)_q, (11001)_q\} \\ &= \text{Orb}(1100)_q \cup \text{Orb}(11100)_q;\end{aligned}$$

If $q = q_{6,p}$ then

$$\begin{aligned}\mathcal{E}(P_{6,p}) &= \{(111000)_q, (011100)_q, (001110)_q, \\ &\quad (000111)_q, (100011)_q, (110001)_q\} \\ &= \text{Orb}(111000)_q.\end{aligned}$$

In Example 3.11 the set of extremal points $\mathcal{E}(P_{n,p})$ is shown to be intimately related to the sequences $(1^{\lfloor n/2 \rfloor} 0^{n - \lfloor n/2 \rfloor})$ and $(1^{\lceil n/2 \rceil} 0^{n - \lceil n/2 \rceil})$ when $n = 3, 5$ and to the sequence $(1^{n/2} 0^{n/2})$ when $n = 4, 6$. We now prove that this is a general result. We set $q = q_{n,p}$ and for $h = 1, \dots, n$ we introduce the vertices

$$(25) \quad \mathbf{v}_{2h-1} := \sigma^h(1^{\lfloor n/2 \rfloor} 0^{n - \lfloor n/2 \rfloor}) = \sum_{k=0}^{\lfloor n/2 \rfloor - 1} q^{k+h-1 \bmod n}$$

$$(26) \quad = \sum_{k \in \underline{K}(h)} q^k$$

where

$$\underline{K}(h) := \{k \in \{0, \dots, n-1\} \mid (k-h+1) \bmod n \leq \lfloor n/2 \rfloor - 1\};$$

$$(27) \quad \mathbf{v}_{2h} := \sigma^h(1^{\lceil n/2 \rceil} 0^{n - \lceil n/2 \rceil}) = \sum_{k=0}^{\lceil n/2 \rceil - 1} q^{k-h+1 \bmod n}$$

$$(28) \quad = \sum_{k \in \overline{K}(h)} q^k$$

where

$$\overline{K}(h) := \{k \in \{0, \dots, n-1\} \mid (k-h+1) \bmod n \leq \lceil n/2 \rceil - 1\}.$$

Remark 3.12. By a direct computation, if n is odd then for every $h = 1, \dots, n$

$$(29) \quad \mathbf{n}_{2h-1} = (-q^{h-2 \bmod n})^\perp$$

$$(30) \quad \mathbf{n}_{2h} = (q^{h-2+\lceil n/2 \rceil \bmod n})^\perp$$

while if n is even then $\mathbf{v}_{2h} = \mathbf{v}_{2h-1}$ hence

$$(31) \quad \mathbf{n}_{2h-1} = \left(1 - \frac{1}{p^{n/2}}\right) (-q^{h-2 \bmod n})^\perp$$

$$(32) \quad \mathbf{n}_{2h} = 0;$$

In view of (29) and of (31), for every n and every h

$$(33) \quad \arg \mathbf{n}_{2h-1} = \left(\frac{(h-2) \bmod n}{n} 2\pi + \frac{\pi}{2} \right) \bmod 2\pi$$

Lemma 3.13. For every $h = 1, \dots, n$

$$(34) \quad k \in \underline{K}(h) \text{ if and only if } q^k \cdot \mathbf{n}_{2h-1} \geq 0;$$

$$(35) \quad k \in \overline{K}(h) \text{ if and only if } q^k \cdot \mathbf{n}_{2h} \geq 0;$$

Proof. For every $k = 0, \dots, n-1$

$$(36) \quad q^k \cdot \mathbf{n}_{2h-1} \geq 0$$

if and only if

$$(37) \quad \cos \left(\arg q^k - \arg \mathbf{n}_{2h-1} \right) = \cos \left(\frac{(k-h+1) \bmod n + 1}{n} 2\pi - \frac{\pi}{2} \right) \geq 0$$

As $(k-h+1) \bmod n \in \{0, \dots, n-1\}$,

$$-\frac{\pi}{2} < \frac{(k-h+1) \bmod n + 1}{n} 2\pi - \frac{\pi}{2} \leq \frac{3\pi}{2},$$

(37) reduces to

$$-\frac{\pi}{2} < \frac{(k-h+1) \bmod n + 1}{n} 2\pi - \frac{\pi}{2} \leq \frac{\pi}{2}$$

Hence (34) follows by the definition of $k \in \underline{K}(h)$ and by recalling that $(k-h+1) \bmod n \in \mathbb{N}$.

To prove (35), we first remark that if n is even then $\mathbf{n}_{2h} = 0$ and (35) is immediate. If otherwise n is odd then

$$\arg \mathbf{n}_{2h} = \left(\frac{(h-2 + \lceil n/2 \rceil) \bmod n}{n} 2\pi + \frac{3\pi}{2} \right) \bmod 2\pi$$

therefore

$$(38) \quad q^k \cdot \mathbf{n}_{2h-1} \geq 0$$

if and only if

$$(39) \quad \cos \left(\frac{(k-h+1) \bmod n + 1 - \lceil n/2 \rceil}{n} 2\pi + \frac{\pi}{2} \right) \geq 0.$$

As $(k-h+1) \bmod n \in \{0, \dots, n-1\}$, for every $k = 0, \dots, n-1$

$$-\frac{\pi}{2} < \frac{(k-h+1) \bmod n + 1}{n} 2\pi - \frac{\pi}{2} \leq \frac{3\pi}{2},$$

(37) reduces to

$$-\frac{\pi}{2} < \frac{(k-h+1) \bmod n + 1}{n} 2\pi - \frac{\pi}{2} \leq \frac{\pi}{2};$$

Relation (35) hence follows by the definition of $\overline{K}(h)$. □

Lemma 3.14. *For every $x \in X_{n,p}$ and for every $h = 1, \dots, n$*

$$(40) \quad (x - \mathbf{v}_{2h-1}) \cdot \mathbf{n}_{2h-1} \leq 0$$

and

$$(41) \quad (x - \mathbf{v}_{2h}) \cdot \mathbf{n}_{2h} \leq 0.$$

Proof. Let

$$x = \sum_{k=0}^{n-1} x_k q^k \in X_{n,p}$$

with $x_0, \dots, x_{n-1} \in \{0, 1\}$. Then for every $h = 1, \dots, n$

$$(x - \mathbf{v}_{2h-1}) \cdot \mathbf{n}_{2h-1} = \sum_{k \in \bar{K}(h)} (x_k - 1)(q^k \cdot \mathbf{n}_{2h-1}) + \sum_{k \notin \bar{K}(h)} x_k (q^k \cdot \mathbf{n}_{2h-1}) \leq 0$$

indeed Lemma 3.13 and $x_k \in \{0, 1\}$ imply that all the terms of the above sums are non-positive. Similarly, again by Lemma 3.13 and by $x_k \in \{0, 1\}$ we may deduce

$$(x - \mathbf{v}_{2h-1}) \cdot \mathbf{n}_{2h-1} = \sum_{k \in \bar{K}(h)} (x_k - 1)(q^k \cdot \mathbf{n}_{2h-1}) + \sum_{k \notin \bar{K}(h)} x_k (q^k \cdot \mathbf{n}_{2h-1}) \leq 0.$$

□

Remark 3.15. *Proposition 2.3 and Lemma 3.14 imply that $X_{n,p}$ is a subset of the polygon whose vertices are $\mathbf{v}_1, \dots, \mathbf{v}_{2n}$, therefore*

$$P_{n,p} = \text{conv}(X_{n,p}) \subseteq \text{conv}\{\mathbf{v}_h \mid h = 1, \dots, n\}.$$

This, together with the fact that $\mathbf{v}_h \in X_{n,p}$ for every $h = 1, \dots, 2n$, implies

$$(42) \quad P_{n,p} = \text{conv}(\{\mathbf{v}_h \mid h = 1, \dots, 2n\}).$$

In particular when n is odd $\mathbf{v}_1, \dots, \mathbf{v}_{2n}$ coincide with the $2n$ vertices of $P_{n,p}$. When n is even, $\mathbf{v}_1 = \mathbf{v}_2, \dots, \mathbf{v}_{2n-1} = \mathbf{v}_{2n}$ are the n vertices of $P_{n,p}$.

Theorem 3.16. *For every $n \geq 1$, $p > 1$ and for every finite alphabet A*

$$(43) \quad \text{conv}(\Lambda_{n,p,A}) = \frac{\max A - \min A}{p^n - 1} \text{conv}(\{\mathbf{v}_h \mid h = 1, \dots, 2n\}) + \sum_{k=0}^{n-1} \min A q^k.$$

Moreover if n is even then

$$(44) \quad \text{conv}(\Lambda_{n,p,A}) = \frac{\max A - \min A}{p^n - 1} \text{conv}(\{\mathbf{v}_{2h} \mid h = 1, \dots, n\}) + \sum_{k=0}^{n-1} \min A q^k.$$

Proof. In view of Remark 3.15, (43) immediately follows by Proposition 3.5. while (44) holds because if n is even then $\mathbf{v}_{2h} = \mathbf{v}_{2h-1}$ for every $h = 1, \dots, n$. □

4. REPRESENTABILITY IN COMPLEX BASE

In this section we give a characterization of the convexity of $\Lambda_{n,p,A}$. To this end we recall some basic elements of the Iterated Function System (IFS) theory. An IFS $\mathcal{F}$ is a finite set of contractive maps over a metric space $\mathbb{X}$, in particular if $\mathbb{X} = \mathbb{C}$ then

$$\mathcal{F} = \{f_i : \mathbb{C} \rightarrow \mathbb{C} \mid i = 1, \dots, m\}$$

for some $m \in \mathbb{N}$ and for every $x, y \in \mathbb{C}$ and $i = 1, \dots, m$

$$|f_i(x) - f_i(y)| \leq c_i |x - y|$$

for some $0 < c_i < 1$. The *Hutchinson operator* acts on the power set of $\mathbb{C}$ as follows

$$\mathcal{F}(X) := \bigcup_{i=1}^m f_i(X) = \bigcup_{i=1}^m \bigcup_{x \in X} f_i(x).$$

By the contraction principle, every IFS has a unique fixed point $R = \mathcal{F}(R)$ that can be constructed starting by any subset of $\mathbb{C}$, indeed for every $X \subset \mathbb{C}$

$$R = \lim_{k \rightarrow \infty} \mathcal{F}^k(X).$$

Example 4.1. If $q \in \mathbb{C}$ and if $|q| > 1$ then

$$\mathcal{F} = \left\{ f_i : x \mapsto \frac{1}{q}(x + a_i) \mid a_i \in \mathbb{R}; i = 1, \dots, m \right\}$$

is an IFS.

We define

$$(45) \quad \mathcal{F}_{q,A} := \left\{ f_i : x \mapsto \frac{1}{q}(x + a_i) \mid a_i \in A \right\}.$$

Lemma 4.2. Let $q \in \mathbb{C}$, $|q| > 1$ and $A = \{a_1, \dots, a_m\} \subset \mathbb{R}$. Then the fixed point of $\mathcal{F}_{q,A}$ is

$$\Lambda_{q,A} := \left\{ \sum_{k=1}^{\infty} \frac{x_k}{q^k} \mid x_k \in A \right\}.$$

Proof. For every

$$x = \sum_{k=1}^{\infty} \frac{x_k}{q^k} \in \Lambda_{q,A}$$

we have

$$f_i(x) = \frac{1}{q}x + \frac{1}{q}a_i = \sum_{k=2}^{\infty} \frac{x_{k-1}}{q^k} + \frac{a_i}{q} \in \Lambda_{q,A}.$$

Moreover if $x_1 = a_i$ then

$$f_i^{-1}(x) = qx - a_i = x_1 + q \sum_{k=2}^{\infty} \frac{x_k}{q^k} - a_i = \sum_{k=1}^{\infty} \frac{x_{k+1}}{q^k} \in \Lambda_{q,A}.$$

Therefore

$$\bigcup_{i=1}^m f_i(\Lambda_{q,A}) = \Lambda_{q,A}$$

namely $\Lambda_{n,p,A}$ is the fixed point of $\mathcal{F}_{q,A}$. □




FIGURE 4. First 6 iterations of $F_{\{0,1\}} = \{x/q_{3,2}, 1/q_{3,2}(x+1)\}$ over $\text{conv}(\Lambda_{3,2,\{0,1\}})$.

If $q = q_{n,p}$ we define

$$\mathcal{F}_{n,p,A} := \mathcal{F}_{q,A}.$$

Lemma 4.3. *For every $n \in \mathbb{N}$, $p > 1$ and $A \subset \mathbb{R}$, with $|A| < \infty$, if $\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$ is convex then*

$$(46) \quad \text{conv}(\Lambda_{n,p,A}) \subseteq \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})).$$

Proof. Consider $\mathbf{w}_1, \dots, \mathbf{w}_{2n}$, the (possibly pairwise coincident) vertices of $\Lambda_{n,p,A}$. By Theorem 3.16, for every $h = 1, \dots, 2n$

$$\begin{aligned}
\mathbf{w}_h &= \frac{\max A - \min A}{p^n - 1} \mathbf{v}_h + \frac{1}{p^n - 1} \sum_{k=0}^{n-1} \min A q^k \\
&= \frac{\max A - \min A}{p^n - 1} \sigma^h(1^l 0^{n-l})_q + \frac{1}{p^n - 1} \sum_{k=0}^{n-1} \min A q^k
\end{aligned}$$

for some $h \in \{1, \dots, n\}$ and $l \in \{\lfloor n/2 \rfloor, \lceil n/2 \rceil\}$. Remark that if $\varepsilon \in \{0, 1\}$ is the last digit of $\sigma^h(1^l 0^{n-l})$ then

$$q\sigma^h(1^l 0^{n-l})_q = \sigma^{h+1}(1^l 0^{n-l})_q + \varepsilon(1 - q^n).$$

Now, as $q^n = p^n$ if $\varepsilon = 0$ then

$$\begin{aligned}
f_1^{-1}(\mathbf{w}_h) &= q\mathbf{w}_h - \min A \\
&= \frac{\max A - \min A}{p^n - 1} (q\sigma^h(1^l 0^{n-l})_q) + \frac{q}{p^n - 1} \sum_{k=0}^{n-1} \min A q^k - \min A \\
&= \frac{\max A - \min A}{p^n - 1} (\sigma^{h+1}(1^l 0^{n-l})_q) + \frac{1}{p^n - 1} \sum_{k=0}^{n-1} \min A q^k \\
&= \mathbf{w}_{h+2}.
\end{aligned}$$

therefore $f_1(\mathbf{w}_{h+2}) = \mathbf{w}_h$. By a similar argument, it is possible to show that if $\varepsilon = 1$ then $f_m(\mathbf{w}_{h+2}) = \mathbf{w}_h$. Thus

$$(47) \quad \mathbf{w}_h \in \mathcal{F}_{q,A}(\{\mathbf{w}_h \mid h = 1, \dots, 2n\})$$

and, by the arbitrariness of $\mathbf{w}_h$,

$$(48) \quad \{\mathbf{w}_h \mid h = 1, \dots, 2n\} \subseteq \mathcal{F}_{q,A}(\{\mathbf{w}_h \mid h = 1, \dots, 2n\})$$

Hence

$$\begin{aligned}
\text{conv}(\Lambda_{n,p,A}) &= \text{conv}(\{\mathbf{w}_h \mid h = 1, \dots, 2n\}) \\
&\subseteq \text{conv}(\mathcal{F}_{q,A}(\{\mathbf{w}_h \mid h = 1, \dots, 2n\})) \\
&= \text{conv}\left(\bigcup_{i=1}^m f_i(\{\mathbf{w}_h \mid h = 1, \dots, 2n\})\right) \\
&\subseteq \text{conv}\left(\bigcup_{i=1}^m f_i(\text{conv}(\{\mathbf{w}_h \mid h = 1, \dots, 2n\}))\right) \\
&= \text{conv}(\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))) \\
&= \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})).
\end{aligned}$$

□

Lemma 4.4. For every $n \in \mathbb{N}$, $p > 1$ and $A \subset \mathbb{R}$

$$(49) \quad \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})) \subseteq \text{conv}(\Lambda_{n,p,A}).$$

Proof. Let $y \in \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$ so that $y = f_i(x)$ for some $i = 1, \dots, m$ and some $x \in \text{conv}(\Lambda_{n,p,A})$. In particular $x = \alpha x_1 + (1 - \alpha)x_2$ for some $x_1, x_2 \in \Lambda_{n,p,A}$ and, consequently,

$$y = \alpha f_i(x_1) + (1 - \alpha)f_i(x_2)$$

because f_i is a linear map. Since $\Lambda_{n,p,q}$ is the fixed point of an IFS containing f_i we have $f_i(x_1), f_i(x_2) \in \Lambda_{n,p,A}$ and, consequently, $y \in \text{conv}(\Lambda_{n,p,A})$. Thesis follows by the arbitrariness of y . $\square$

Lemma 4.5. *For every $n \in \mathbb{N}$, $p > 1$ and $A \subset \mathbb{R}$, $\Lambda_{n,p,A}$ is convex if and only if $\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$ is convex.*

Proof. If $\Lambda_{n,p,A}$ is convex, then the convexity of $\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$ follows by

$$(50) \quad \text{conv}(\Lambda_{n,p,A}) = \Lambda_{n,p,A} = \mathcal{F}_{n,p,A}(\Lambda_{n,p,A}) = \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})).$$

If otherwise $\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$ is convex, then by Lemma 4.3

$$\text{conv}(\Lambda_{n,p,A}) \subseteq \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$$

while Lemma 4.4 implies

$$\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})) \subseteq \text{conv}(\Lambda_{n,p,A});$$

therefore

$$\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})) = \text{conv}(\Lambda_{n,p,A});$$

and by the uniqueness of the fixed point of $\mathcal{F}_{n,p,A}$

$$\text{conv}(\Lambda_{n,p,A}) = \Lambda_{n,p,A}.$$

$\square$

Theorem 4.6. *The set of representable numbers in base $q_{n,p}$ and alphabet $A = \{a_i \mid i = 1, \dots, m\}$ is convex if and only if*

$$(51) \quad \max_{i=1, \dots, m-1} a_{i+1} - a_i \leq \frac{\max A - \min A}{p^n - 1}.$$

Proof. By applying Proposition 2.9 with $P = P_{n,p}$ and

$$t_i = a_i \frac{p^n - 1}{\max A - \min A},$$

we get that (51) holds if and only if

$$\bigcup_{i=1}^m (P_{n,p} + t_i)$$

is convex. Then (51) is equivalent to the convexity of

$$\begin{aligned} \bar{P} &:= \frac{\max A - \min A}{q(p^n - 1)} \bigcup_{i=1}^m (P_{n,p} + t_i) + \frac{1}{q(p^n - 1)} \sum_{k=0}^{n-1} \min A q^k. \\ &= \bigcup_{i=1}^m \frac{1}{q} \left(\frac{\max A - \min A}{q(p^n - 1)} P_{n,p} + \frac{1}{q(p^n - 1)} \sum_{k=0}^{n-1} \min A q^k + a_i \right) \end{aligned}$$




FIGURE 5. $\Lambda_{3,2^{1/3}+0.1k,\{0,1\}}$, with $k = 0, \dots, 5$, is approximated with the set of expansions with length 14.

By Theorem 3.16 we have

$$\bar{P} = \bigcup_{i=1}^m \frac{1}{q} (\text{conv}(\Lambda_{n,p,A}) + a_i) = \mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A})),$$

therefore $\mathcal{F}_{n,p,A}(\text{conv}(\Lambda_{n,p,A}))$ is convex if and only if (51) holds. This hence follows by Lemma 4.5. $\square$

Corollary 4.7. *Let $A = \{a_i \mid i = 1, \dots, m\}$. If*

$$\max_{i=1, \dots, m-1} a_{i+1} - a_i \leq \frac{\max A - \min A}{p^n - 1}$$

then every

$$x \in \frac{\max A - \min A}{p^n - 1} P_{n,p} + \sum_{k=0}^{n-1} \min A q_{n,p}^k$$

has a representation in base $q_{n,p}$ and alphabet A .

Proof. It immediately follows by Theorem 4.6. $\square$

Example 4.8. If $p = 2^{1/n}$ and $A = \{0, 1\}$ then $\Lambda_{n,p,A}$ is a convex set coinciding with $P_{n,p}$. In particular $\Lambda_{n,p,A}$ is an $2n$ -gon if n is odd and it is an n -gon if n is even.

Example 4.9. If $A = \{0, 1, \dots, \lfloor p^n \rfloor\}$ then $\Lambda_{n,p,A}$ is a convex set or, equivalently,

$$\frac{\lfloor p^n \rfloor}{p^n - 1} P_{n,p}$$

is completely representable.

REFERENCES

- [1] S. Akiyama and J. M. Thuswaldner. A survey on topological properties of tiles related to number systems. *Geom. Dedicata*, 109:89–105, 2004.
- [2] Z. Daróczy and I. Kátai. Generalized number systems in the complex plane. *Acta Math. Hungar.*, 51(3-4):409–416, 1988.
- [3] V. A. Dimitrov, G. A. Jullien, and Miller W.C. Complexity and fast algorithms for multiexponentiation. 31:141–147, 1985.
- [4] Ch. Frougny and A. Surarerks. On-line multiplication in real and complex base. *Computer Arithmetic, IEEE Symposium on*, 0:212, 2003.
- [5] W. J. Gilbert. Geometry of radix representations. In *The geometric vein*, pages 129–139. Springer, New York, 1981.
- [6] W. J. Gilbert. Arithmetic in complex bases. *Math. Mag.*, 57(2):77–81, 1984.
- [7] W. J. Gilbert. Complex bases and fractal similarity. *Ann. Sci. Math. Québec*, 11(1):65–77, 1987.
- [8] editor Heckbert, P. S. *Graphics Gems IV*. Academic Press, 1994.
- [9] I. Kátai and B. Kovács. Canonical number systems in imaginary quadratic fields. *Acta Math. Acad. Sci. Hungar.*, 37(1-3):159–164, 1981.
- [10] I. Kátai and J. Szabó. Canonical number systems for complex integers. *Acta Sci. Math. (Szeged)*, 37(3-4):255–260, 1975.
- [11] D. E. Knuth. An imaginary number system. *Comm. ACM*, 3:245–247, 1960.
- [12] D. E. Knuth. *The art of computer programming. Vol. 2*. Addison-Wesley Publishing Co., Reading, Mass., first edition, 1971. Seminumerical algorithms, Addison-Wesley Series in Computer Science and Information Processing.
- [13] V. Komornik and P. Loreti. Expansions in complex bases. *Canad. Math. Bull.*, 50(3):399–408, 2007.
- [14] W. Penney. A “binary” system for complex numbers. *J. ACM*, 12(2):247–248, 1965.
- [15] D. Piché. Complex bases, number systems and their application to the fractal wavelet image coding. *PhD thesis (Univ. Waterloo, Ontario)*, 0, 2002.
- [16] J. Pineda. A parallel algorithm for polygon rasterization. 22(4):17–20, 1988.
- [17] J. A. Solinas. Efficient arithmetic on Koblitz curves. *Des. Codes Cryptogr.*, 19(2-3):195–249, 2000. Towards a quarter-century of public key cryptography.

DIPARTIMENTO DI SCIENZE DI BASE E APPLICATE PER L’INGEGNERIA, SAPIENZA UNIVERSITÀ DI ROMA